Importance of first trimester scan – 2 of 4

Twin pregnancy

Tiran Dias¹

Sri Lanka Journal of Obstetrics and Gynaecology 2011; 33: 72-76

Introduction

Twin pregnancies have a greater incidence of perinatal morbidity and mortality, primarily attributed to the uteroplacental insufficiency, premature births and antepartum complications¹. Monochorionic (MC) twins account for 20% of all spontaneous twin pregnancies, but yet carry five time higher perinatal morbidities and mortalities than dichorionic (DC) twins²⁻³. This is due to the presence of placental vascular anastomoses allowing interfetal transfusion, which accounts for the 15-20% of twin-twin transfusion syndrome (TTTS) and a further 25% of severe birthweight discordances⁴. Moreover, most of the complications in monochorionic twins are early onset (before 20 weeks) and if not treated majority of them will die before 25 weeks. Additionally, in the event of a single-twin intrauterine death (IUD), the acute intertwin transfusion poses a more profound risk for the surviving MC twin than for the DC twin⁵. Consequences for the surviving co-twin include severe neurological impairment, death and premature delivery⁵. Therefore it is vital to determine the chorionicity in the first trimester.

Determination of chorionicity

Chorionicity can be accurately determined during first trimester ultrasonography. Monochorionic twin pregnancy can be diagnosed in the presence of the T-sign and dichorionic with the λ -sign or when two separate placental masses are present⁶⁻⁷ (Figures 1 and 2). The λ -sign was first described by Bessis and Papernik (1981), and referred to the triangular tissue projection extending from the base of the inter-twin membrane in early dichorionic pregnancies with a single fused placenta⁸. The term λ -sign has been used interchangeably with the 'twin peak' sign⁹. The absence of the λ -sign leads to the finding of the T-sign characteristic of monochorionicity. Largest study of the accuracy of the T/ λ -signs and number of placentas

in the first trimester diagnosis of chorionicity has shown that chorionicity determination is feasible and very accurate at 11 to 14 weeks gestation⁷. Dias *et al* in that study reported the sensitivity of T/ λ -signs and number of placentas for the detection of monochorionicity was 100% with only one dichorionic pregnancy incorrectly assigned as monochorionic⁷.

Figure 1. 'T' sign in monochorionic twin pregnancy.

Figure 2. Lambda sign (λ -sign) in dichorionic twin pregnancy.

¹ Senior Lecturer in Obstetrics and Gynaecology, Department of Obstetrics and Gynaecology, Faculty of Medicine, University of Colombo, Sri Lanka.

E-mail: thiran_dias@yahoo.com

Figure 3. Two different placental mass in dichorionic twins.

As the pregnancy advances into the second trimester, regression of the chorion frondosum leads to the gradual loss of the λ -sign. This change explains the observation that the λ -sign is not visible in about 7% of dichorionic pregnancies by 20 weeks' gestation¹⁰. Therefore, it is mandatory to assign chorionicity in the first trimester. If there is uncertainty about the chorionicity, thermal image of the membrane attachment to the placenta should be retained in the case notes and a second opinion should be sought or the woman should be referred to a specialist without delay, as chorionicity is best determined before 14 weeks (RCOG)¹¹.

Twin pregnancy dating

Routine dating of pregnancy from a first trimester crown-rump length (CRL) is superior to the use of menstrual dates¹²⁻¹⁴. Accurate dating of twin pregnancies is vital as twin pregnancies are at increased risk of perinatal morbidity and mortality compared to singletons, mainly as a consequence of both preterm delivery and fetal growth restriction.

There are two main concerns in twin pregnancy dating. Firstly, whether CRL charts derived from singleton pregnancies can be used to accurately date twin pregnancy. Secondly, which twin's measurement should be considered for dating? Dias *et al* concluded that singleton CRL charts can be used reliably to date twin pregnancy with the difference of 1-2 days (Table 1)¹⁵. Furthermore, larger twin's CRL is more pragmatic for dating as pathological largeness in early pregnancy is not biologically plausible but smallness¹⁵. After 14 weeks as for singletons, head circumference can be reliably used for twin pregnancy dating in second trimester up to 25 weeks¹⁹.

Twin pregnancy orientation

A reproducible method for antenatal labeling of twins is important in the management of all twin pregnancies. This is to ensure that biometry from longitudinal growth scans are consistently allocated to the same twin at each visit. Inconsistent allocation of twin in each visit may result in the 'yo-yoing' of fetal growth as smaller and larger twin sizes are swapped repeatedly during the course of the pregnancy. Additionally, when screening for aneuploidies is undertaken, there must be a reliable and accurate system in place to ensure that invasive prenatal diagnosis or selective fetal reduction is carried out on the at-risk or affected twin, respectively¹¹. Each fetus within the twin pregnancy can be orientated at the 11-14 week ultrasound assessment, the fetus contained in the gestational sac closest to the maternal cervix is designated as twin one^{20,21}. The relative orientation of the fetuses to each other (Figures 4 and 5) is then defined as either lateral (left/right) or vertical (top/bottom). Lateral fetal orientation is associated with an inter-twin membrane running vertically along the longitudinal axis of the uterus and a vertical fetal orientation is associated with an inter-twin membrane running horizontally across the longitudinal axis of the uterus (Figures 4 and 5).

Table 1. The mean difference between observed (from ultrasound) and expected (from IVF history) measurements expressed in days with 95% confidence interval in parentheses¹⁵ (Dias *et al*)

CRL formulae	Mean difference in gestation (days)		
	Singleton	Bigger twin	Smaller twin
Robinson ¹⁶	1.41 days (1.15 to 1.68)	2.4 days (2.4 to 2.6)	0.91 days (0.55 to 1.30)
Rossavik ¹⁷	0.14 days (0.01 to 0.28)	1.27 days (1.05 to 1.5)	-0.51 days (-0.30 to -0.72)
Von Kaisenberg ¹⁸	-0.54 days (-0.41 to -0.67)	0.58 days (0.36 to 0.8)	-1.18 days (-0.97 to -1.4)

Figure 4. Diagrammatic representation of twin orientation relative to the longitudinal axis of the uterus - left/right orientation.

Figure 5. Diagrammatic representation of twin orientation relative to the longitudinal axis of the uterus - top/bottom orientation.

Table 2. Risk of trisomy 21 at 12 weeks

Age (yrs)	Risk for trisomy 21 at 12 weeks		
	Singleton	Monozygotic	Dizygotic
20	1100	1100	550
25	1000	1000	500
30	650	650	325
31	550	550	275
32	450	450	225
33	400	400	200
34	300	300	150
35	250	250	125
36	200	200	100
37	150	150	75
38	120	120	60
39	90	90	45
40	70	70	35
41	50	50	25
42	40	40	20

Aneuploidy screening in twin pregnancy

Risk of underlying chromosomal problems in each fetus depends on zygosity. In dizygotic twins the maternal age-related risk for chromosomal abnormalities for each fetus is the same as in singleton pregnancies. As majority of dichorionic twins are dizygotic the chance that at least one fetus is affected by a chromosomal defect is twice as high as in singleton pregnancies. Since all the monochorionic twins are monozygotic their risk for a chromosomal abnormality affecting both fetuses is the same as in singleton pregnancies (Table 2).

Risk assessment for chromosomal abnormalities in twin pregnancies can be effectively done by a combination of maternal age, fetal NT thickness, FHR and maternal serum free β -hCG and PAPP-A²²⁻²³. Most importantly standardized values of serum biochemistry should be used in the risk calculation formula in twin pregnancy²³.

In dichorionic twins the fetus specific risk for trisomy 21 is determined by the differences in NT. The detection rate (>90%) and false positive rate (3% per fetus or 6% per pregnancy) are similar to those in singleton pregnancies²².

Increased NT in at least one of the fetuses could be an early manifestation of TTTS in Monochorionic twins²⁴. Therefore, the false positive rate of screening (8% per pregnancy) is higher than in DC twins. As all monochorionic twins are monozygotic, they invariably concordant for fetal karyotype. In monochorionic twins the risk for trisomy²¹ is calculated for each fetus, based on maternal age, fetal NT and maternal serum biochemistry and then the average risk between the two fetuses is considered to be the risk for the pregnancy as a whole²⁵.

Amniocentesis in twins is effective in providing a reliable karyotype for both fetuses and the procedure-related fetal loss rate is about 2%. In the case of chorionic villous sampling, the procedure-related fetal loss rate is also about 2%, but in about 1% of cases there may be a diagnostic error, either due to sampling the same placenta twice or cross-contamination.

In pregnancies discordant for chromosomal defects the main options are either selective fetocide or expectant management. Selective fetocide after 16 weeks of gestation is associated with three-fold increase in risk of spontaneous abortion compared to reduction before 16 weeks. Consequently, if the parents request invasive testing it is preferable to perform CVS rather than amniocentesis.

References

1. Sherer DM. Adverse perinatal outcome of twin pregnancies according to chorionicity: review of the literature. *American Journal of Perinatology* 2001; **18**: 23-37.
2. Dias T, Bhide A, Thilaganathan B. Early pregnancy growth and pregnancy outcome in twin pregnancies. *Ceylon Med J* 2010; **55**(3): 80-4.
3. Acosta-Rojas R, Becker J, Munoz-Abellana B, et al. Twin chorionicity and the risk of adverse perinatal outcome. *International Journal of Obstetrics and Gynaecology* 2007; **96**: 98-102.
4. Barigye O, Pasquini L, Galea P, et al. High risk of unexpected late fetal death in monochorionic twins despite intensive ultrasound surveillance: a cohort study. *PLoS Medicine* 2005; **2**(6): e172.
5. Ong SS, Zamora J, Khan KS, Kilby MD. Prognosis for the co-twin following single-twin death: a systematic review. *BJOG* 2006; **113**: 992-8.
6. Sepulveda W, Sebire NJ, Hughes K, Odibo A, Nicholaides KH. The lambda sign at 10-14 weeks of gestation as a predictor of chorionicity in twin pregnancies. *Ultrasound Obstet Gynecol* 1996; **7**: 421-3.
7. Dias T, Arcangeli T, Bhide A, Napolitano R, Mahsud-Dornan S, Thilaganathan B. First-trimester ultrasound determination of chorionicity in twin pregnancy. *Ultrasound Obstet Gynecol* 2011 Jan 31. doi: 10.1002/uog.8956.
8. Bessis R, Papernik E. Echographic imagery of amniotic membranes in twin pregnancies. *Twin Research 3: Twin Biology and Multiple Pregnancy*. Alan R Liss: New York; 1981: 183-7.
9. Finberg HJ. The 'twin peak' sign: reliable evidence of dichorionic twinning. *J Ultrasound Med* 1992; **11**: 571-7.
10. Sepulveda W. Chorionicity determination in twin pregnancies: double trouble. *Ultrasound Obstet Gynecol* 1997; **10**: 79-81.
11. Royal College of Obstetricians and Gynaecologists. Green-top Guideline No. 51. Management of monochorionic twin pregnancy, December 2008.
12. Neilson JP. Ultrasound for fetal assessment in early pregnancy. *Cochrane Database of Systematic Reviews* 2001, Issue 2. Art. No.: CD000182. DOI: 10.1002/14651858.CD000182.
13. Taipale P, Hiilesmaa V. Predicting delivery date by ultrasound and last menstrual period in early gestation. *Obstet Gynecol* 2001; **97**: 189-94.
14. Mongelli M, Wilcox M, Gardosi J. Estimating the date of confinement. *Am J Obstet Gynecol* 1996; **174**: 278-81.
15. Dias T, Mahsud-Dornan S, Thilaganathan B, Papageorgiou A, Bhide A. First-trimester ultrasound dating of twin pregnancy: are singleton charts reliable? *BJOG* 2010; **117**(8): 979-84. Epub 2010 May 19.
16. Robinson HP, Fleming JEE. A critical evaluation of sonar "crown-rump length" measurements. *BJOG* 1975; **82**: 702-10.
17. Von Kaisenberg CS, Fritzer E, Kuhling H, Jonat W. Fetal transabdominal biometry at 11-14 weeks of gestation. *Ultrasound Obstet Gynecol* 2002; **20**: 564-74.
18. Rossavik IK, Torjusen GO, Gibbons WE. Conceptual age and ultrasound measurements of gestational sac and crown-rump length in in vitro fertilization pregnancies. *Fertil Steril* 1988; **49**: 1012-7.

19. Dias T, Arcangeli T, Bhide A, Mahsud-Dornan S, Papageorghiou A, Thilaganathan B. Second-trimester assessment of gestational age in twins: validation of singleton biometry charts. *Ultrasound Obstet Gynecol* 2011; **37**(1): 34-7.
20. Chudleigh T, Thilaganathan B. First trimester ultrasound. In *Obstetric Ultrasound: How, Why and When* (3rd edn), Elsevier, Churchill Livingstone: London, 2004; 29-49.
21. Dias T, Ladd S, Mahsud-Dornan S, Bhide A, Papageorghiou AT, Thilaganathan B. Systematic labelling of twin pregnancies on ultrasound: limitations and guidance. *Ultrasound Obstet Gynecol* 2011 Mar 11. doi: 10.1002/uog.8990. [Epub ahead of print]
22. Wright D, Syngelaki A, Staboulidou I, Cruz Jde J, Nicolaides KH. Screening for trisomies in dichorionic twins by measurement of fetal nuchal translucency thickness according to the mixture model. *Prenat Diagn* 2011; **31**(1): 16-21. doi: 10.1002/pd.2643.
23. Spencer K, Kagan KO, Nicolaides KH. Screening for trisomy 21 in twin pregnancies in the first trimester: an update of the impact of chorionicity on maternal serum markers. *Prenat Diagn* 2008; **28**(1): 49-52.
24. Kagan KO, Gazzoni A, Sepulveda-Gonzalez G, Sotiriadis A, Nicolaides KH. Discordance in nuchal translucency thickness in the prediction of severe twin-to-twin transfusion syndrome. *Ultrasound Obstet Gynecol* 2007; **29**(5): 527-32.
25. Cleary-Goldman J, Berkowitz RL. First trimester screening for Down syndrome in multiple pregnancy. *Semin Perinatol* 2005; **29**(6): 395-400.

Subscription Information

ISSN 1391-7536

Subscription rates

Individual copies	-	Rs 300/=
Subscription for one year	-	Rs 1000/=

Please make cheques payable to
"Sri Lanka College of Obstetricians and Gynaecologists"

For information related to the Journal please contact:

SLCOG office
 No 112, Model Farm Road
 Colombo 8
 Sri Lanka.

Manuscripts can be submitted on E-mail to:
deepalsenaka@yahoo.co.uk

Instructions to authors can be downloaded
 from the Journal website:
www.slcof.lk